



HAL
open science

A multilevel approach to individual and organizational predictors of stress and fatigue among healthcare workers of a university hospital: A longitudinal study

Oumou Daouda, René Bun, Karim Bouziad, Katuska Miliani, Anastasia Essa-Eworo, Florence Espinasse, Delphine Seytre, Anne Casetta, Simone Nérome, Adelaide Nascimento, et al.

► To cite this version:

Oumou Daouda, René Bun, Karim Bouziad, Katuska Miliani, Anastasia Essa-Eworo, et al.. A multilevel approach to individual and organizational predictors of stress and fatigue among healthcare workers of a university hospital: A longitudinal study. *Occupational and Environmental Medicine*, BMJ Publishing Group, 2022, 10.1101/2022.01.24.22269481 . hal-03587531v2

HAL Id: hal-03587531

<https://hal-cnam.archives-ouvertes.fr/hal-03587531v2>

Submitted on 10 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



OPEN ACCESS

Original research

Multilevel approach to individual and organisational predictors of stress and fatigue among healthcare workers of a university hospital: a longitudinal study

Oumou Salama Daouda ¹, René Sosata Bun,² Karim Ait Bouziad,¹ Katuska Miliiani,³ Anastasia Essa-Eworo,³ Florence Espinasse,⁴ Delphine Seytre,⁵ Anne Casetta,⁶ Simone Nérôme,⁷ Adelaide Nascimento,⁸ Pascal Astagneau,^{2,3} Laura Temime,¹ Mounia N Hocine¹

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/oemed-2022-108220>).

¹MESuRS Laboratory, Conservatoire National des Arts et Metiers, Paris, Île-de-France, France

²IPLESP, INSERM, Sorbonne Université, Paris, Île-de-France, France

³CPIAS Ile de France, Paris, Île-de-France, France

⁴Hôpital Ambroise Paré, AP-HP, Paris, Île-de-France, France

⁵Hôpital Avicenne, AP-HP, Paris, Île-de-France, France

⁶Hôpital Cochin, AP-HP, Paris, Île-de-France, France

⁷Hôpital Beaujon, AP-HP, Paris, Île-de-France, France

⁸CRTD Laboratory, Conservatoire National des Arts et Metiers, Paris, Île-de-France, France

Correspondence to

Dr Oumou Salama Daouda, MESuRS Laboratory, Conservatoire National des Arts et Metiers, Paris, France; oumou-salama.daouda@lecnam.net

OSD and RSB contributed equally.

Received 10 January 2022
Accepted 22 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Daouda OS, Bun RS, Ait Bouziad K, et al. *Occup Environ Med Epub ahead of print*: [please include Day/Month/Year]. doi:10.1136/oemed-2022-108220

ABSTRACT

Objective Healthcare workers (HCWs) are at high risk of experiencing stress and fatigue due to the demands of their work within hospitals. Improving their physical and mental health and, in turn, the quality and safety of care requires considering factors at both individual and organisational/ward levels. Using a multicentre prospective cohort, this study aims to identify the individual and organisational predictors of stress and fatigue of HCWs in several wards from university hospitals.

Methods Our cohort consists of 695 HCWs from 32 hospital wards drawn at random within four volunteer hospital centres in Paris-area. Three-level longitudinal analyses, accounting for repeated measures (level 1) across participants (level 2) nested within wards (level 3) and adjusted for relevant fixed and time-varying confounders, were performed.

Results At baseline, the sample was composed by 384 registered nurses, 300 auxiliary nurses and 11 midwives. According to the three-level longitudinal models, some predictors were found in common for both stress and fatigue (low social support from supervisors, work overcommitment, sickness presenteeism and number of beds per ward). However, specific predictors for high level of stress (negative life events, low social support from colleagues and breaks frequently cancelled due to work overload) and fatigue (longer commuting duration, frequent use of interim staff in the ward) were also found.

Conclusion Our results may help identify at-risk HCWs and wards, where interventions to reduce stress and fatigue should be focused. These interventions could include manager training to favour better staff support and overall safety culture of HCWs.

INTRODUCTION

There is a growing research interest about stress in healthcare workers (HCWs), as the prevalence of nurses affected by negative mental states is high.^{1,2} A recent meta-analysis including 45 539 nurses worldwide in 49 countries across multiple specialties estimated 11.2% prevalence of burnout among global nurses.³

There is a vast literature exploring occupational predictors of both stress and fatigue among HCWs,

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Healthcare workers (HCWs) have high levels of perceived stress and fatigue, particularly in medical fields highly exposed to infectious risks.
- ⇒ Previous studies show that the high physical and mental demands, as well as unfavourable work environment and organisation, can increase stress and fatigue levels of HCWs.
- ⇒ However, there is a lack of studies exploring the complex dynamics of the link between working conditions and HCW health, using multilevel design (HCWs nested within wards), including ward/organisational level predictors.

WHAT THIS STUDY ADDS

- ⇒ The ward-level environment significantly influences the stress and fatigue of HCWs in addition to individual factors and time variations.
- ⇒ Supervisors providing low social support and with low safety culture, work overcommitment, sickness presenteeism and working in smaller wards were identified as predictors of both high stress and fatigue of HCWs.
- ⇒ Negative life events (whether occurring in personal or in professional life), low support from colleagues and breaks frequently cancelled due to work overload are specific predictors of high level of stress. While commuting duration, frequent use of interim staff and working in a medical ward were associated with high level of fatigue of HCWs.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ In this study, we can identify some areas for improvement to better prevent stress and fatigue for HCWs. High stress and fatigue can be reduced through mutual and specific organisational intervention strategies.

including studies with cross-sectional, longitudinal and experimental designs. Low social support from supervisors and colleagues, as well as younger age and being female, was found to be predictors of

stress among HCWs.^{4–6} In addition, several studies highlight the importance of job satisfaction and favourable work environment in order to prevent stress of HCWs. Indeed, unacceptable work schedule, work overcommitment and shift work appeared to lead to higher stress of HCWs.^{7–9} Also, being highly committed to the organisation was associated with both role and work overload and work-related stress.⁹

Similarly, several studies explored the relationship between HCWs fatigue and some aspects of working conditions such as low social support from supervisor and colleagues, older age as well as carrying for both children and elders.^{10–12} Psychological job demands and unfavourable work environment (including shift work organisation and work schedule characteristics) also emerged as significant predictors of fatigue among HCWs.^{10–13} In addition, HCWs with higher levels of overcommitment were more prone to experience chronic fatigue syndrome.¹⁴

As HCWs are responsible for playing their role in keeping patient safety, observing safety principles while delivering daily care can lead to a decrease in adverse events and damages.¹⁵ Negative HCW perception of patient safety can also have a detrimental effect on stress and fatigue, as suggested by recent studies reporting negative correlations between patient safety culture and the level of stress and fatigue.^{16–17}

HCWs fatigue and mental health are important elements to monitor for continued HCWs well-being and patient safety.¹⁸ Stress and fatigue in the workplace negatively impact productivity and increase absenteeism of HCWs¹⁹ and may result in non-optimal quality of care for patients in healthcare settings.²⁰ For instance in 2020, the average absenteeism rate in French public hospitals was estimated 9.5% among a representative sample of 300 hospitals²¹ versus 5.1% nation wide (<https://www.ayming.fr/wp-content/uploads/sites/11/2020/09/Barometre-absenteisme-2020-Ayming.pdf>).

Still, a better and more global understanding of how individual-level and organisational-level factors interacts with stress and fatigue of HCWs in hospitals is needed, especially in the French context where studies on hospital staff stress and fatigue are scarce. The originality of our study comes from its multilevel design. We designed the study longitudinally to observe the possible evolutions of fatigue and stress regarding the organisational factors that may vary over time in order to better identify the levers of organisational control that could reduce stress and fatigue.

Here, we propose a three-level longitudinal analysis using individual and organisational/ward factors related to stress and fatigue of HCWs, using a 1-year longitudinal data collected from 730 individuals in 32 French hospital wards.

MATERIAL AND METHODS

Study design and data collection

We designed a multicentre study on the individual and organisational predictors of stress and fatigue among HCWs at the Hospitals of Paris, the STRIPPS (*Stress au Travail et Risque Infectieux chez les Patients et le Personnel Soignant*) study.²² The study was conducted between February 2018 and July 2019 and data were collected on midwives, registered and auxiliary nurses. An auxiliary nurse, also known as a healthcare assistant, works closely with healthcare professionals (nurses and physicians). He/she performs duties such as washing and dressing patients, or making beds, in healthcare facilities. HCWs were recruited from four voluntary French University general care hospitals. Eight wards per participating hospital were drawn at random among wards employing at least 30 HCWs.

Data were collected longitudinally every 4 months during 1 year by two different interviewers for all included participants as follows: t0, corresponding to the first collection during the HCWs inclusion visit; t1, t2, t3, corresponding to follow-up visits at 4 months, 8 months and 12 months. For the first data collection (t0), dates and times of visits by an interviewer were drawn randomly for each participating ward. For later data collections, individual appointments were made with each included HCWs. Data were collected through questionnaire-based interviews at both levels (ward and individual levels).

Individual-level variables

To collect individual-level variables, HCWs of included wards were interviewed individually at 4 times of visits (t0, t1, t2 and t3). The individual questionnaire collected demographic and occupational characteristics, including age, sex, professional status, contractual situation, experience in the hospital and daily working hours. Individual data related to work organisation, including schedule, nightshift, extra hours, mealtimes and rest periods, were also collected. Social support from colleagues and from supervisor was also assessed using the social support dimension of the Job Content Questionnaire (JCQ), which was developed and validated by Karasek *et al.*²³ Social support dimension includes eight 4-points Likert scale items (from 1 to 4), with a resulting score ranging from 8 to 32 (from very low to very high social support). In addition, work overcommitment was measured using the work overcommitment dimension of the Effort-Reward Imbalance questionnaire (ERI).²⁴ This dimension includes six 4-points Likert scale items (from 1 to 4), with a resulting score ranging from 6 to 24. Low total score indicates low work overcommitment. The reliability statistics results showed that Cronbach's alphas for social support and work overcommitment were 0.79 and 0.76, respectively, indicating acceptable scale reliability. Finally, the management support for patient safety dimension was measured using the French version of the hospital survey on patient safety culture.²⁵

Ward-level variables

To collect data for each of the 32 wards, the hospital health executives (nurse managers) were interviewed at t0 only regarding the following characteristics: medical specialty, number of beds per ward, proportion of double rooms, frequency of tasks performed outside the ward, patient/physician ratio, patient/paramedics ratio and use of external healthcare service providers (ie, interim staff).

Outcome variables

Two primary outcomes were considered:

1. Perceived stress, assessed with the Perceived Stress Scale 10-item scale (PSS-10), was developed by Cohen *et al.*²⁶ and validated in French.²⁷ The PSS-10 questionnaire includes 10 5-point Likert items (from 0 to 4), with a resulting score ranging from 0 to 40 (from very low to very high perceived stress). The reliability statistics results showed that Cronbach's alpha for PSS-10 scale was 0.85, indicating acceptable scale reliability.
2. Fatigue, assessed with the Pichot Fatigue Scale.²⁸ The Pichot Scale includes eight 4-point Likert items (from 1 to 4), with a score ranging from 8 to 32 (from very low to very high fatigue). Cronbach's alpha for Pichot scale was 0.91, indicating acceptable scale reliability.

Why study stress and fatigue together

Although stress and fatigue are distinct phenomena, both can be considered as mediating factors of behavioural consequences,

including absenteeism, turnover and productivity loss of HCWs.^{29 30} Earlier studies considered both stress and fatigue together as outcomes.³¹ Here, we assumed that both may result from the high physical and mental demands to which HCWs are exposed, with possible consequences in terms of patient safety. Identifying factors associated with both high stress and fatigue may, therefore, help improve HCW physical and mental health and patient care.

Missing data imputation

For missing data, multiple imputations were performed on validated questionnaire items only (JCQ, PSS-10, Pichot and ERI questionnaires). Data were imputed using multiple imputation, using the R *mice* package.³² The *mice* package allows performing imputation of continuous and categorical variables in a context of multilevel and longitudinal data. For all questionnaire items included in the imputation model, missing data were assumed to be missing at random.

Statistical analysis and modelling

We conducted analyses to identify predictors of stress and fatigue levels of participating HCWs. First, in order to validate the use of a three-level longitudinal model, we built two unconditional models (ie, null models, with no independent variables) with two levels (ie, time and individual levels) and three levels (ie, time, individual and ward levels), for each outcome. Indeed, before conducting multivariate multilevel models, performing null models is strongly encouraged.²⁹ We then compared the two unconditional models using the Akaike's Information Criterion (AIC).³⁰ Lower AIC for three-level unconditional models validated the using of three-level models to predict stress and fatigue. In addition, using null models, intraclass correlations were computed, in order to quantify how much response variable variance is shared, across different combinations of levels.

Due to the large number of explanatory variables and to avoid collinearity with some of them, we proceeded to two steps of selection. Specifically:

1. First, univariate analyses were performed for all individual-level variables in order to reduce the number of variables to include in the multivariate model. Then, any variables with univariate associations with p values ≤ 0.20 were included in the multivariate model. This cut-off was chosen to exclude variables of questionable importance.³³ Despite the fact that the study was longitudinal, univariate analysis was performed at baseline. This technique is often used and allows to capture the maximum of information on the variables and the outcome.³⁴ Indeed, the response rate at the baseline is naturally higher than at follow-ups.
2. Second, we developed a three-level multivariate model on each outcome, using the AIC for variable selection, using only variable retained in the first step. All data analyses were conducted using the R software.³⁵

RESULTS

Response rate

Overall, the response rate for all included HCWs who answered the questionnaires at the four visits was 73.5% (510 out of 694), corresponding to 2040 responses in total for all visits. In total, 695 HCWs were included at t0, 644 at t1, 578 at t2 and 556 at t3, with an overall of 2473 observations. A general response rate at t0 is impossible to compute due to the voluntary participation of HCWs present at the time of inclusion in the study. However, at the time of inclusion, the participation rate based

on the total number of staff working in the participating wards was $695/2473=28.1\%$. In addition, the mean number of observations per ward decreases across time points. We observed 22.8 observations at t0, 21.25 at t1, 19.1 at t2 and 18.5 at t3.

An auxiliary nurse failed to answer any of the four questions of social support from supervisor of the Karasek questionnaire and was, thus, excluded from all analyses.

HCW characteristics

The final study sample consisted of 694 HCWs as follows: registered nurses ($n=384$) (55.3%), auxiliary nurses ($n=299$) (43.1%) and midwives ($n=11$) (1.6%). Overall, the female/male gender ratio was 5.5, with 588 (84.7%) female respondents. The majority of HCWs were permanent staff members ($n=616$) (88.9%) compared with temporary ($n=58$) (8.4%) and contractual ($n=19$) (2.7%) staff members. The average number of years of experience was 9 years (9.6), and more than a half of the respondents had supervising responsibilities ($n=365$) (52.6%) (table 1).

Ward characteristics

In total, 32 wards were included from various medical fields as follows: 14 (43.8%) in surgery and obstetrics, 11 (34.4%) in 'other medical specialty wards' and 7 (21.9%) in intensive care unit (ICU) and reanimation. The average number of beds per ward was 35.5 (SD=18.5), and the proportion of double rooms was approximately 20% (table 2).

In participating wards, the average patient/physician ratio was 2.9 (4.0), whereas the patient/paramedics ratio was 0.8 (0.3). In the vast majority of participating wards, work was organised in 3 to 8-hour shifts, while 16% of wards worked in 2 to 12-hour shifts.

Mean-level change across 1 year of survey

The trajectories for each participant across the total sample from baseline to the last time point are presented in online supplemental table 1 for time-varying individual-level factors. Significant differences among the four times of visits were observed for schedule assignment frequency, number of nightshifts on duty over the last months, irregularity of mealtimes, number of breaks cancelled due to work overload, number of visits to the occupational safety and health department, sickness presenteeism and social support from supervisor.

Outcome characteristics

The distribution of PSS-10 (stress) and Pichot (fatigue) scores in the whole sample are presented in figure 1, respectively, in (A) and (B). The overall mean score is equal to 16.5 (7.0) out of 40 points for stress, and 11.0 (7.9) out of 32 points for fatigue. For fatigue only, we observed an increasing trend ($p=0.028$) of the means across the time of visits (online supplemental table 2). For both stress and fatigue, significant differences were observed between the four hospitals ($p<0.001$) (online supplemental table 3). Figure 2A describes the trend of stress and fatigue level across time of visits and hospital. Additional figures on the distribution of stress and fatigue by hospital are available at online supplemental figures 1 and 2, respectively.

According to unconditional models, AICs from three-level unconditional models are lower than those from two-level unconditional models for both stress and fatigue (online supplemental table 4). Hence, the results obtained validate the use of the three-level models to analyse both outcomes. Additional information on intraclass correlations is available under online

Table 1 Characteristics of individuals at the time of inclusion t0

	Total (N=694)
Age (years)	
≤25	125 (18.0%)
26–35	236 (34.0%)
36–45	143 (20.6%)
46–55	126 (18.2%)
>55	64 (9.2%)
Gender	
Female	588 (84.7%)
Male	106 (15.3%)
Profession	
Registered nurses	384 (55.3%)
Auxiliary nurse (AS)	299 (43.1%)
Midwives	11 (1.6%)
Years of work in the hospital	
Mean (SD)	9.0 (9.6)
Professional status	
Temporary/contractual CDD	58 (8.4%)
Contract/contractual CDI	19 (2.7%)
Permanent/titulaire	616 (88.9%)
Missing	1
Supervising responsibility	
No	329 (47.4%)
Yes	365 (52.6%)
Weekly working hours	
Mean (SD)	36.8 (4.2)
Knowledge of the schedule	
Never	13 (1.9%)
Sometimes	74 (10.7%)
Always	607 (87.5%)
Participation to the schedule	
Never	321 (46.3%)
Sometimes	266 (38.3%)
Always	107 (15.4%)
48 hours weekly rest	
No	295 (42.6%)
Yes	398 (57.4%)
Missing	1
Number of breaks per day	
Mean (SD)	1.8 (1.0)
Total duration of breaks per day (min)	
Mean (SD)	15.3 (9.3)
Transportation	
Personal car	357 (51.4%)
Public transportation	286 (41.2%)
Other (walking, bike, motorbike)	51 (7.3%)
Commuting time to work	
<1 hour	290 (41.8%)
1 hour–2 hours	305 (43.9%)
>2 hours	99 (14.3%)
HSOPSC item-1*	
Agree	213 (30.7%)
Neutral	106 (15.3%)
Do not agree	375 (54.0%)
HSOPSC item-8†	
Agree	220 (31.7%)
Neutral	96 (13.9%)

continued

Table 1 continued

	Total (N=694)
Do not agree	377 (54.4%)
Missing	1
HSOPSC item-9‡	
Agree	465 (67.2%)
Neutral	92 (13.3%)
Do not agree	135 (19.5%)
Missing	2
HSOPSC item-10§	
Agree	405 (58.4%)
Neutral	116 (16.7%)
Do not agree	172 (24.8%)
Missing	1
Support from colleagues	
Mean (SD)	13.2 (2.0)
Support from hierarchy	
Mean (SD)	11.7 (2.7)
Work overcommitment	
Mean (SD)	15.5 (2.7)
*HSOPSC item-1: from the Hospital Survey on Patient Safety Culture. 'Hospital management provides a work climate that promotes patient safety'.	
†HSOPSC item-8: from the Hospital Survey on Patient Safety Culture. 'The actions of hospital management show that patient safety is a top priority'.	
‡HSOPSC item-9: from the Hospital Survey on Patient Safety Culture. 'Hospital management seems interested in patient safety only after an adverse event happens'.	
§HSOPSC item-10: from the Hospital Survey on Patient Safety Culture. 'Hospital units work well together to provide the best care for patients'.	

supplemental table 4). A summary of missing values according to the four validated scales, by time of visits, is available in online supplemental table 5).

The best three-level multivariate models identified after model selection using AIC are described in table 3 along with model fit measures.

Individual and organisational-level predictors of stress

At individual level, negative life events (whether occurring in personal or in professional life), frequent cancellation of breaks, low social support from supervisor and colleagues, low perception of hospital management attitude towards patient safety culture, work overcommitment, as well as sickness presenteeism, were significantly associated to high perceived stress of HCWs. At ward-level, medical specialty (with less stress in ICU than in surgical and obstetrical units or wards with other medical specialties), and number of beds (with more stress in smaller wards) emerged as significant predictors of perceived stress.

Individual and organisational-level predictors of fatigue

At individual level, gender (with more fatigue in female participants), longer commuting time to work, low social support from supervisor, low perception of hospital management attitude towards patient safety culture, work overcommitment, as well as sickness presenteeism, emerged as significant predictors of high level of fatigue of HCWs. Significant predictors of fatigue at ward level were medical specialty (with more fatigue in wards with other medical specialties), higher rates of interim use and number of beds in the ward (with more fatigue in smaller wards). High fatigue level was also associated to time-level variable (time of visit, with an increasing trend over 12-month period).

Table 2 Characteristics of included wards

	Total (N=32)
Specialty	
Surgery/obstetrics	14 (43.8%)
Other medical specialties*	11 (34.4%)
ICU/reanimation	7 (21.9%)
Number of beds	
Mean (SD)	35.5 (18.5)
Proportion of double bedrooms	
Mean (SD)	0.2 (0.2)
Going outside the ward	
Never	1 (3.2%)
Sometimes	11 (35.5%)
Fairly often	14 (45.2%)
Always	5 (16.1%)
Missing	1
Ratio patient/physician	
Mean (SD)	2.9 (4.0)
Use of external healthcare services	
Never	8 (25.0%)
Sometimes	17 (53.1%)
Fairly often	7 (21.9%)
Ratio patient/paramedics	
Mean (SD)	0.8 (0.3)
Time schedule	
2*12 hours	5 (15.6%)
3*8 hour	27 (84.4%)

*Other medical specialties: cardiology, geriatrics, gastroenterology, infectious diseases, internal medicine, nephrology, oncology, pulmonology, rheumatology, urology.
ICU, intensive care unit.

DISCUSSION

Main findings

In this longitudinal study, we identified various individual and organisational risk factors of stress and fatigue of HCWs. In particular, a supervisor providing low social support, low safety culture, work overcommitment, sickness presenteeism and working in smaller wards were identified as predictors of both high stress and fatigue.

In addition, breaks frequently cancelled due to work overload, negative life events (whether occurring in personal or in professional life), low social support from colleagues and working in non-surgical/obstetrical wards are specific predictors of high stress. However, longer commuting duration and frequent use of interim staff were associated with high level of fatigue.

Comparison with the literature

Many of our findings are consistent with those reported in previous studies investigating the determinants of stress or fatigue of HCWs.

In particular, the influence of the low social support from supervisor on both stress and fatigue was underlined in an earlier French study.³¹ Consistently with our results, this same study also showed higher fatigue in small to medium hospital wards, and in work environments where staff frequently had to go outside the ward as well as lower energy levels and more frequent sleep difficulties when use of interim staff was frequent.³¹ Safety climate perceptions were found to be significantly related to HCW stress and fatigue in several recent studies.³⁶

Moreover, our finding that work overcommitment and sickness presenteeism, another indicator of overcommitment, were significant predictors of stress and fatigue, is supported by a recent French study in which overcommitment was found to favour emotional exhaustion and increase the risk of burnout in French HCWs.³⁷

However, some other factors including younger age, being female, shift work as well as unacceptable work schedule were previously reported in the literature as associated with stress, but not in our results.^{5 7 38} Regarding fatigue, factors previously reported in the literature but not found in our results were support from colleagues, and work schedule characteristics such as total working hours working, overtime number of monthly night (if at least half of the working hours are between midnight and 8:00) and evening (if more than half of the shift hours are between the hours of 16:00 and midnight) shifts and shift length (12 hours vs 8 hours).³¹ This latter could be explained by a potential lack of power due to the moderate sample size, there were a few wards with 12-hour shifts compared with those with an 8-hour shifts.

Interestingly, our results showed some factors associated with stress and/or fatigue not previously reported in the literature. Regarding stress, associated factors were negative life events, breaks cancelled due to work overload, in line with role overload, which a well-known factor of stress. Regarding fatigue, associated factors were longer commuting duration and frequent use of interim staff in the ward. Indeed, temporary contracts require dedicated time for staff training for managers, leading to fatigue. Another factor related to stress and fatigue is the number of beds (higher stress and fatigue in smaller wards). This could be explained by the fact that strain of working on understaffed wards is harder to manage in smaller wards. In addition, smaller wards are associated to lower level of management and HCWs staff, who are required to perform a wide variety of tasks.

Strengths

First, one strength of this study is its longitudinal nature. A few studies explored stress and fatigue longitudinally; however, the majority of currently available studies are cross-sectional. Furthermore, the high response rate of wards and HCWs, as well as the large sample size and the inclusion of wards of different size and activity, represent strength of this study. In addition, the large panel of sociodemographics, health and occupational characteristics of HCWs collected over time allow performing robust and well-adjusted multivariate analysis.

Second, stress and fatigue were explored together. To our knowledge, no previous study had proposed a single model to identify factors associated with high levels of combined stress and fatigue while accounting for time in hospital setting. In fact, a previous study conducted in French ICU attempted to predict stress and fatigue using demographic and occupational was based on cross-sectional survey.³¹

Finally, the power of the model used, which takes into account the complexity of the data, namely, the longitudinal design and the multiwards collection of the data. In recent years, these types of models have been frequently used³⁹ considering the idea that longitudinal data could be analysed at three levels of nesting (eg, repeated measures (level 1), collected across individuals (level 2) and within different wards (level 3)).

Limitations

However, our study has some limitations. First, only four hospitals <https://www.sciencedirect.com/topics/>




Figure 1 Distribution of PSS-10 and Pichot scores, respectively in (A) and (B) among the whole sample. The vertical dotted lines represent the mean of the PSS-10 and Pichot scores. PSS-10, Perceived Stress Scale 10-item scale.

nursing-and-health-professions/tertiary-care-center in Paris-area were included, which is not representative for other areas of France. In fact, only services in public Paris-area hospitals were included in the study, so results may not be generalisable to wards in private hospitals or outside the Paris region, Future

studies including HCWs and more hospitals from other cities in France are needed to verify the results generated in this study.

Second, due to the in-person interview, a risk of bias could be present due to the discomfort from having to reply face-to-face to some sensitive questions; however, ensuring anonymity of the




Figure 2 PSS-10 and Pichot scores means, respectively in (A) and (B), by hospital and time of visits. PSS-10, Perceived Stress Scale 10-item scale.

Table 3 Final three-level models for outcomes (perceived stress and fatigue) using a stepwise approach with AIC criterion

	Stress (PSS-10) (n=2422)*			Fatigue (Pichot) (n=2431)*		
	Estimates	CI	P value	Estimates	CI	P value
Fixed effects†						
Time	-0.00	-0.16 to 0.16	0.957	0.48	0.31 to 0.66	<0.001
Interviewer (ref=1 (for hospitals A and C))						
2 (for hospitals B and D)	1.62	0.85 to 2.40	<0.001	-0.36	-1.47 to 0.76	0.531
<i>HCW-level variables</i>						
Gender (ref=female)						
Male	-0.83	-1.73 to 0.06	0.068	-1.52	-2.63 to -0.40	0.008
Age (ref=<25)						
26–35				1.26	0.09 to 2.44	0.035
36–45				0.50	-0.79 to 1.78	0.450
46–55				-1.01	-2.37 to 0.35	0.147
>55				-2.83	-4.49 to -1.17	0.001
Commuting time to work (ref=>1 hour)						
>2 hour				1.82	0.80 to 2.83	<0.001
1 hour–2 hours				0.30	-0.37 to 0.96	0.382
Personal life event (ref=no)						
Yes, negative	2.27	1.81 to 2.74	<0.001			
Yes, positive	0.46	-0.19 to 1.12	0.164			
Professional life event (ref=no)						
Yes, negative	1.62	1.11 to 2.12	<0.001			
Yes, positive	0.40	-0.34 to 1.14	0.289			
Break cancelled due to work overload (ref=never)						
Almost never	1.04	0.27 to 1.82	0.008			
Quite often	1.54	0.77 to 2.32	<0.001			
Very often	1.78	0.92 to 2.65	<0.001			
Professional status (ref=temporary CDD)						
Contract (contractuel CDI)				2.23	-0.53 to 4.99	0.114
Permanent (titulaire)				0.45	-1.04 to 1.93	0.554
Knowing work schedule in advance (ref=never)						
Sometimes	-2.30	-4.81 to 0.22	0.074			
Always	-1.87	-4.23 to 0.49	0.121			
Support from colleagues—Karasek subscale						
Support from supervisor—Karasek subscale	-0.12	-0.24 to -0.01	0.035			
HSOPSC item-1‡ (ref=agree)	-0.23	-0.33 to -0.13	<0.001	-0.34	-0.45 to -0.24	<0.001
HSOPSC item-8§ (ref=agree)						
Neutral	0.00	-1.01 to 1.02	0.995	0.70	-0.52 to 1.91	0.260
Not agree	0.81	-0.10 to 1.71	0.082	1.76	0.81 to 2.70	<0.001
Work overcommitment—Siegrist subscale						
Sickness presenteeism (ref=never)	0.71	0.63 to 0.78	<0.001	0.65	0.56 to 0.73	<0.001
Sickness presenteeism (ref=never)						
Almost never	0.04	-0.48 to 0.57	0.871	0.63	0.04 to 1.22	0.037
Quite often	1.04	0.45 to 1.63	0.001	2.48	1.82 to 3.15	<0.001
Very often	1.61	0.55 to 2.66	0.003	4.03	2.84 to 5.22	<0.001
<i>Ward-level variables</i>						
Specialty (ref=surgery/obstetrics)						
Other medical specialties¶	-0.36	-1.30 to 0.59	0.459	1.59	0.24 to 2.94	0.021
ICU/reanimation	-1.08	-2.01 to -0.15	0.023	-0.24	-1.60 to 1.13	0.733
Number of beds per ward	-0.03	-0.05 to -0.01	0.009	-0.04	-0.07 to -0.00	0.024
Going outside the ward (ref=never)						
Sometimes	1.28	-0.87 to 3.43	0.243	0.98	-2.09 to 4.04	0.533
Often	1.53	-0.70 to 3.76	0.178	0.86	-2.30 to 4.03	0.592
Always	1.87	-0.44 to 4.18	0.112	2.59	-0.70 to 5.89	0.123
Use of interim nurses (ref=never)						
Sometimes	0.05	-0.81 to 0.91	0.915	-0.36	-1.59 to 0.88	0.571

continued

Table 3 continued

	Stress (PSS-10) (n=2422)*			Fatigue (Pichot) (n=2431)*		
	Estimates	CI	P value	Estimates	CI	P value
Often	0.96	-0.14 to 2.06	0.088	1.96	0.35 to 3.56	0.017
Random effects σ^2 (σ)*						
Level 1—time	17.80 (4.23)			22.04 (4.69)		
Level 2—healthcare worker	12.30 (3.51)			19.69 (4.44)		
Level 3—ward	0.00 (0.00)			0.50 (0.70)		
Marginal R ² /Conditional R ² **	0.507/NA			0.299/0.634		
AIC	14 743.16			15 433.57		

*The slightly different samples for each outcome refer to the difference between the factors included in the final model for each outcome, following the variable selection. Indeed, the number of missing values per variable has an impact on the number of observations that will be considered in the final mode.

†In multilevel models, fixed effects are usually equivalent to the regression coefficients, while random effects usually account for the underlying structure of the data and characterised by estimates of variability σ^2 (σ). Fixed effects can be interpreted as slopes in the traditional sense.

‡HSOPSC item-1: item-1 from the Hospital Survey on Patient Safety Culture. 'Hospital management provides a work climate that promotes patient safety'.

§HSOPSC item-8: item-8 from the Hospital Survey on Patient Safety Culture. 'The actions of hospital management show that patient safety is a top priority'.

¶The marginal R² considers only the variance of the fixed effects, while the conditional R² takes both the fixed and random effects into account.

AIC, Akaike information criterion; CDD, Fixed-term contract; CDI, Permanent employment contract; HCW, healthcare worker; ICC, intra-class correlation coefficient; ICU, intensive care unit; PSS-10, Perceived Stress Scale 10-item scale.

participants was used to minimise such bias. Another possible source of bias in data collection is the presence of two different interviewers assigned to two hospitals each. However, we were able to consider this bias as we included this interviewer-related variable into the multilevel model. In the stress model, there was a significant interviewer effect, with higher stress in hospitals B and D. However, interviewer effect was not significant in the fatigue model.

Third, we were not able to investigate stress and fatigue outcomes for the physicians given low response rates, as questions regarding work organisation were less adequate than for nurses. Therefore, physicians were excluded from our sample of HCWs.

Insights for designing potential interventions

From these models, we can identify some areas for improvement to better prevent stress and fatigue of HCWs: (1) perception of the hierarchy (low social support from supervisor, low perception of hospital management attitude towards patient safety culture), (2) work overcommitment and (3) sickness presenteeism. Breaks frequently cancelled due to work overload and low social support from colleagues were also found as significant as specific predictors for stress level. For fatigue specifically, long commuting duration and use of external staff are also identified as predictors. Mutual and specific preventive programmes for reducing stress and fatigue of HCWs could be implemented in order to reduce this burden, targeting on the most at-risk groups.

CONCLUSION

This research question is important given the influence on quality of patient care of high stress work environments.¹⁸ Our results may (1) help identify at-risk HCWs and wards, where interventions to reduce stress and fatigue could be focused. (2) These interventions could include manager training to favour better staff support and overall safety culture among HCWs.

Acknowledgements The sponsor was Assistance Publique—Hôpitaux de Paris (Délégation à la Recherche Clinique et à l'Innovation). The authors would like to thank Zahoua Tina Habbal, the AP-HP and all participating wards and HCWs for their time and their contribution to this study.

Contributors OSD is the study guarantor. OSD: methodology, software, formal analysis, writing—original draft, writing—review and editing, visualisation, validation; RSB: methodology, formal analysis, writing—original draft,

writing—review and editing, visualisation, validation; KAB: methodology, formal analysis, writing—review and editing, visualisation, validation; KM: conceptualisation, writing—review and editing, visualisation, validation; AE-E: investigation. FE: conceptualisation, supervision, project administration; DS: conceptualisation, supervision, project administration; AC: conceptualisation, supervision, project administration; SN: project administration, supervision, project administration; AN: conceptualisation, visualisation, funding acquisition, validation; PA: conceptualisation, methodology, formal analysis, writing—review and editing, visualisation, supervision, funding acquisition, validation; LT: conceptualisation, methodology, formal analysis, writing—review and editing, visualisation, supervision, funding acquisition, validation; MNH: conceptualisation, methodology, formal analysis, writing—review and editing, visualisation, supervision, funding acquisition, validation.

Funding The study was publicly funded through the Research on the Evaluation of the Healthcare System Performance Programme, Ministry of Health, France (grant number PREPS-16-261).

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval The study protocol was elaborated in collaboration with the AP-HP Department of medical policy and the Department of care and of paramedical activities and was approved after presentation to the Directorate General and the Committee on hygiene, safety and working conditions. It obtained both an agreement from the French Committee for the Protection of Persons (CPP) on 11/14/2017 and clearance from the French Data Protection Authority (CNIL) on 12/14/2017 (IDRCB N° 2017-A02939-44). Potential participants were informed of the study through an information letter. Verbal consent was obtained by the interviewer at the beginning of each interview. Participants were guaranteed confidentiality and anonymity of responses. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. The data that support the findings of this study are available from the authors, upon reasonable request. It can be addressed to Pr Pascal Astagneau (pascal.astagneau@aphp.fr), the project leader.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is

properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Oumou Salama Daouda <http://orcid.org/0000-0001-9682-1599>

REFERENCES

- Maharaj S, Lees T, Lal S. Prevalence and risk factors of depression, anxiety, and stress in a cohort of Australian nurses. *Int J Environ Res Public Health* 2018;16:61.
- Durand A-C, Bompard C, Sportiello J, et al. Stress and burnout among professionals working in the emergency department in a French university hospital: prevalence and associated factors. *Work* 2019;63:57–67.
- Woo T, Ho R, Tang A, et al. Global prevalence of burnout symptoms among nurses: a systematic review and meta-analysis. *J Psychiatr Res* 2020;123:9–20.
- Assis BBde, Azevedo C, Moura CdeC, et al. Factors associated with stress, anxiety and depression in nursing professionals in the hospital context. *Rev Bras Enferm* 2022;75Suppl 3:e20210263.
- Chatzigianni D, Tsounis A, Markopoulos N, et al. Occupational stress experienced by nurses working in a Greek regional Hospital: a cross-sectional study. *Iran J Nurs Midwifery Res* 2018;23:450–7.
- El-Hneiti M, Shaheen AM, Bani Salameh A, et al. Predictors of nurses' stress working with older people admitted to acute care setting. *Int J Older People Nurs* 2019;14:e12222.
- Tripodi D, Roedlich C, Laheux MA, et al. Stress perception among employees in a French university hospital. *Occup Med* 2012;62:216–9.
- Kath LM, Stichter JF, Ehrhart MG, et al. Predictors and outcomes of nurse leader job stress experienced by AWHONN members. *J Obstet Gynecol Neonatal Nurs* 2013;42:E12–25.
- Bolino MC, Turnley WH. The personal costs of citizenship behavior: the relationship between individual initiative and role overload, job stress, and work-family conflict. *J Appl Psychol* 2005;90:740–8.
- Jalilian H, Shouroki FK, Azmoon H, et al. Relationship between job stress and fatigue based on job Demand-control-support model in hospital nurses. *Int J Prev Med* 2019;10:56.
- Chen J, Daraiseh NM, Davis KG, et al. Sources of work-related acute fatigue in United States Hospital nurses. *Nurs Health Sci* 2014;16:19–25.
- Scott LD, Hwang W-T, Rogers AE. The impact of multiple care giving roles on fatigue, stress, and work performance among hospital staff nurses. *J Nurs Adm* 2006;36:86–95.
- Han K, Trinkoff AM, Geiger-Brown J. Factors associated with work-related fatigue and recovery in hospital nurses working 12-hour shifts. *Workplace Health Saf* 2014;62:409–14.
- Li M, Shu Q, Huang H, et al. Associations of occupational stress, workplace violence, and organizational support on chronic fatigue syndrome among nurses. *J Adv Nurs* 2020;76:1151–61.
- HM M, H H, K H. Managers And Nurses Function Of Safe Patients' Care From The Nurses Perspective. *JHPM* 2012;1:7–14.
- Asefzadeh S, Kalhor R, Tir M. Patient safety culture and job stress among nurses in Mazandaran, Iran. *Electron Physician* 2017;9:6010–6.
- Al Ma'mari Q, Sharour LA, Al Omari O. Fatigue, burnout, work environment, workload and perceived patient safety culture among critical care nurses. *Br J Nurs* 2020;29:28–34.
- Jang HJ, Kim O, Kim S, et al. Factors affecting physical and mental fatigue among female Hospital nurses: the Korea nurses' health study. *Healthcare* 2021;9:201.
- Mollazadeh M, Saraei M, Mehrdad R, et al. Sickness absenteeism of healthcare workers in a teaching hospital. *Hosp Pract Res* 2018;3:6–10.
- Ruotsalainen S, Jantunen S, Sinervo T. Which factors are related to Finnish home care workers' job satisfaction, stress, psychological distress and perceived quality of care? - a mixed method study. *BMC Health Serv Res* 2020;20:896.
- Enquête FHF : les ressources humaines face la seconde vague Covid-19 - Fédération Hospitalière de France (FHF). Available: <https://www.fhf.fr/Presse-Communication/Actualites/Enquete-FHF-les-ressources-humaines-face-a-la-seconde-vague-Covid-19> [Accessed 17 Sep 2021].
- Stress at Work and Infectious Risk in Patients and Caregivers - Full Text View - ClinicalTrials.gov. Available: <https://clinicaltrials.gov/ct2/show/NCT03532321> [Accessed 2 Sep 2021].
- Karasek R, Brisson C, Kawakami N, et al. The job content questionnaire (JCQ): an instrument for internationally comparative assessments of psychosocial job characteristics. *J Occup Health Psychol* 1998;3:322–55.
- Siegrist J, Starke D, Chandola T, et al. The measurement of effort-reward imbalance at work: European comparisons. *Soc Sci Med* 2004;58:1483–99.
- Occelli P, Quenon J-L, Kret M, et al. Validation of the French version of the hospital survey on patient safety culture questionnaire. *Int J Qual Health Care* 2013;25:459–68.
- Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983;24:385–96.
- Lesage F-X, Berjot S, Deschamps F. Psychometric properties of the French versions of the perceived stress scale. *Int J Occup Med Environ Health* 2012;25:178–84.
- Pichot P, Brun JP. [Brief self-evaluation questionnaire for depressive, asthenic and anxious dimensions]. *Ann Med Psychol* 1984;142:862–5.
- Monsalves MJ, Bangdiwala AS, Thabane A, et al. LEVEL (Logical Explanations & Visualizations of Estimates in Linear mixed models): recommendations for reporting multilevel data and analyses. *BMC Med Res Methodol* 2020;20:3.
- Akaike H. A new look at the statistical model identification. *IEEE Trans Automat Contr* 1974;19:716–23.
- Jones G, Hocine M, Salomon J, et al. Demographic and occupational predictors of stress and fatigue in French intensive-care registered nurses and nurses' aides: a cross-sectional study. *Int J Nurs Stud* 2015;52:250–9.
- van BS, Groothuis-Oudshoorn K, Vink G. Mice: multivariate imputation by Chained equations, 2021. Available: <https://CRAN.R-project.org/package=mice> [Accessed 8 Sep 2021].
- Kim EM, Li G, Kim M. Development of a risk score to predict postoperative delirium in patients with hip fracture. *Anesth Analg* 2020;130:79–86.
- Sekine L, Arns B, Fabro BR, et al. Convalescent plasma for COVID-19 in hospitalised patients: an open-label, randomised clinical trial. *Eur Respir J* 2022;59:2101471.
- R: the R project for statistical computing. Available: <https://www.r-project.org/> [Accessed 12 Nov 2021].
- McGhan GE, Ludlow NC, Rathert C, et al. Variations in workplace safety climate perceptions and outcomes across healthcare provider positions. *Journal of Healthcare Management* 2020;65:202–15.
- Sérole C, Auclair C, Prunet D, et al. The forgotten health-care occupations at risk of Burnout-A burnout, job Demand-Control-Support, and Effort-Reward imbalance survey. *J Occup Environ Med* 2021;63:e416–25.
- Muhamad Robot R, Mohd Fauzi MF, Mat Saruan NA, et al. Why so stressed? A comparative study on stressors and stress between hospital and non-hospital nurses. *BMC Nurs* 2021;20:2.
- Peugh JL, Heck RH. Conducting three-level longitudinal analyses. *J Early Adolesc* 2017;37:7–58.

Supplemental materials

A multilevel approach to individual and organizational predictors of stress and fatigue among healthcare workers of a university hospital: a longitudinal study

by Oumou Salama Daouda¹, René Sosata Bun, Karim Ait Bouziad, Katiuska Miliiani, Anastasia Essaworo, Florence Espinasse, Delphine Seytre, Anne Casetta, Simone Nérome, Adelaide Nascimento, Pascal Astagneau, Laura Temime, Mounia N Hocine

¹ Correspondence concerning this article should be addressed to Ms. Oumou S. Daouda, MESuRS laboratory, Cnam, 292 rue Saint Martin, 75003 Paris, France (email: oumou-salama.daouda@lecnam.net)

Table of contents

Supplementary figure 1: Distribution of PSS-10 score. by hospital. The vertical dashed lines represent the means of the PSS-10 scores by hospital

Supplementary figure 2: Distribution of Pichot score by hospital. The vertical dashed lines represent the means of the Pichot scores by hospital

Supplementary table 1. Characteristics of individuals and missing values at the times of visits (t0, t1, t2 and t3)


Supplementary table 2: Means and ranges of outcomes variables (PSS-10 and Pichot scores), by time of visits

Supplementary table 3: Outcomes (PSS-10 and Pichot scores), and missing values by hospital


Supplementary table 4: Unconditional 2 and 3-level models for outcomes of perceived stress and fatigue

Supplementary table 5: Summary of missing values according to validated scales, by time of visits

Supplementary figure 1: Distribution of PSS-10 score by hospital. The vertical dashed lines represent the means of the PSS-10 scores by hospital



Supplementary figure 2: Distribution of Pichot score by hospital. The vertical dashed lines represent the means of the Pichot scores by hospital



Supplementary table 1. Characteristics of individuals and missing values at the times of visits (t0, t1, t2 and t3)

Two-sided ANOVA tests were performed for continuous variables and Chi square tests were performed for qualitative variables

	t0 (N=694)	t1 (N=644)	t2 (N=578)	t3 (N=556)	Total (N=2472)	p value
Work schedule of last months						0.636
Daily	458 (66.0%)	436 (67.7%)	381 (65.9%)	357 (64.2%)	1632 (66.0%)	
Nightly	198 (28.5%)	185 (28.7%)	170 (29.4%)	168 (30.2%)	721 (29.2%)	
Day and Night	38 (5.5%)	23 (3.6%)	27 (4.7%)	31 (5.6%)	119 (4.8%)	
Schedule assignment frequency						< 0.001
Mostly	76 (11.0%)	47 (7.3%)	23 (4.0%)	36 (6.5%)	182 (7.4%)	
Always	618 (89.0%)	597 (92.7%)	555 (96.0%)	519 (93.5%)	2289 (92.6%)	
Missing	0	0	0	1	1	
Nightshift/duty on last months						< 0.001
No	564 (81.4%)	380 (59.1%)	324 (56.2%)	319 (57.7%)	1587 (64.4%)	
Yes	129 (18.6%)	263 (40.9%)	252 (43.8%)	234 (42.3%)	878 (35.6%)	
Missing	1	1	2	3	7	
Number of nightshift/duties						< 0.001
Mean (SD)	0.6 (1.7)	1.3 (2.2)	1.5 (2.3)	1.5 (2.4)	1.2 (2.2)	
Missing	4	5	2	3	14	
Work schedule variation						0.762
Never	404 (58.2%)	360 (55.9%)	334 (57.8%)	319 (57.4%)	1417 (57.3%)	
Fairly often	122 (17.6%)	123 (19.1%)	94 (16.3%)	100 (18.0%)	439 (17.8%)	
Almost Never	146 (21.0%)	132 (20.5%)	133 (23.0%)	114 (20.5%)	525 (21.2%)	
Very often	22 (3.2%)	29 (4.5%)	17 (2.9%)	23 (4.1%)	91 (3.7%)	
Overtime hours						0.098
Never	238 (34.3%)	209 (32.5%)	177 (30.6%)	179 (32.2%)	803 (32.5%)	
Fairly often	205 (29.5%)	209 (32.5%)	208 (36.0%)	189 (34.0%)	811 (32.8%)	
Almost Never	173 (24.9%)	180 (28.0%)	149 (25.8%)	139 (25.0%)	641 (25.9%)	

Very often	78 (11.2%)	46 (7.1%)	44 (7.6%)	49 (8.8%)	217 (8.8%)	
Irregularity of meal time						0.047
Never	41 (5.9%)	51 (7.9%)	34 (5.9%)	37 (6.7%)	163 (6.6%)	
Fairly often	191 (27.6%)	182 (28.3%)	188 (32.5%)	175 (31.5%)	736 (29.8%)	
Almost Never	62 (8.9%)	55 (8.5%)	63 (10.9%)	69 (12.4%)	249 (10.1%)	
Very often	399 (57.6%)	356 (55.3%)	293 (50.7%)	275 (49.5%)	1323 (53.5%)	
Missing	1	0	0	0	1	
Number of canceled breaks						< 0.001
Never	48 (6.9%)	75 (11.6%)	67 (11.6%)	65 (11.7%)	255 (10.3%)	
Fairly often	279 (40.2%)	266 (41.3%)	229 (39.6%)	221 (39.7%)	995 (40.3%)	
Almost Never	129 (18.6%)	128 (19.9%)	152 (26.3%)	162 (29.1%)	571 (23.1%)	
Very often	238 (34.3%)	175 (27.2%)	130 (22.5%)	108 (19.4%)	651 (26.3%)	
Number of visits to the Occupational safety and health (OSH) department						< 0.001
Mean (SD)	0.3 (0.5)	0.2 (0.5)	0.1 (0.4)	0.2 (0.5)	0.2 (0.5)	
Personal life events						0.148
No	408 (58.8%)	383 (59.6%)	350 (60.6%)	337 (60.6%)	1478 (59.8%)	
Yes, negative	217 (31.3%)	192 (29.9%)	166 (28.7%)	141 (25.4%)	716 (29.0%)	
Yes, positive	69 (9.9%)	68 (10.6%)	62 (10.7%)	78 (14.0%)	277 (11.2%)	
Missing	0	1	0	0	1	
Professional life events						0.557
No	488 (70.5%)	463 (71.9%)	390 (67.6%)	381 (68.6%)	1722 (69.8%)	
Yes, negative	151 (21.8%)	139 (21.6%)	137 (23.7%)	123 (22.2%)	550 (22.3%)	
Yes, positive	53 (7.7%)	42 (6.5%)	50 (8.7%)	51 (9.2%)	196 (7.9%)	
Missing	2	0	1	1	4	
Sickness presenteeism						0.005
Never	159 (22.9%)	192 (29.9%)	122 (21.1%)	154 (27.7%)	627 (25.4%)	
Fairly often	218 (31.4%)	178 (27.7%)	174 (30.2%)	148 (26.7%)	718 (29.1%)	
Almost Never	275 (39.6%)	242 (37.7%)	251 (43.5%)	235 (42.3%)	1003 (40.6%)	
Very often	42 (6.1%)	30 (4.7%)	30 (5.2%)	18 (3.2%)	120 (4.9%)	
Missing	0	2	1	1	4	
Marital status						0.371

Couple	382 (55.0%)	360 (55.9%)	336 (58.3%)	328 (59.4%)	1406 (57.0%)	
Single	312 (45.0%)	284 (44.1%)	240 (41.7%)	224 (40.6%)	1060 (43.0%)	
Missing	0	0	2	4	6	
Commuting time to work						0.109
<1 h	290 (41.8%)	270 (41.9%)	253 (43.8%)	243 (43.7%)	1056 (42.7%)	
>2h	99 (14.3%)	91 (14.1%)	54 (9.3%)	60 (10.8%)	304 (12.3%)	
1-2 h	305 (43.9%)	283 (43.9%)	271 (46.9%)	253 (45.5%)	1112 (45.0%)	
Support from colleagues						0.476
Mean (SD)	13.2 (2.0)	13.1 (2.0)	13.1 (2.0)	13.0 (2.0)	13.1 (2.0)	
Support from hierarchy						< 0.001
Mean (SD)	11.7 (2.7)	11.5 (2.5)	11.4 (2.6)	11.1 (2.8)	11.4 (2.7)	
Work overcommitment						0.558
Mean (SD)	15.5 (2.7)	15.4 (2.7)	15.3 (2.6)	15.3 (2.6)	15.4 (2.7)	

Supplementary table 2: Means and ranges of outcomes variables (PSS-10 and Pichot scores), by time of visits

	t0 (N=694)	t1 (N=644)	t2 (N=578)	t3 (N=556)	Total (N=2472)	p_value
PSS-10 score						0.126
Mean (SD)	17.0 (7.0)	16.5 (7.0)	16.3 (7.0)	16.2 (7.1)	16.5 (7.0)	
Range	0.0 - 38.0	0.0 - 38.0	1.0 - 40.0	0.0 - 39.0	0.0 - 40.0	
Pichot score						0.028
Mean (SD)	10.4 (7.8)	11.5 (8.0)	10.9 (7.9)	11.4 (7.9)	11.0 (7.9)	
Range	0.0 - 32.0	0.0 - 32.0	0.0 - 32.0	0.0 - 32.0	0.0 - 32.0	

Supplementary table 3: Outcomes (PSS-10 and Pichot scores), and missing values by hospital

	A (N=610)	B (N=538)	C (N=801)	D (N=523)	Total (N=2472)	p_value
PSS-10 score						< 0.001
Mean (SD)	16.0 (7.2)	17.0 (7.2)	15.3 (6.8)	18.6 (6.6)	16.5 (7.0)	
Missing	4	12	2	4	22	
Pichot score						< 0.001
Mean (SD)	11.4 (8.0)	10.3 (7.9)	10.1 (7.3)	12.8 (8.3)	11.0 (7.9)	
Missing	0	0	2	1	3	

Supplementary table 4: Unconditional 2 and 3-level models for outcomes of perceived stress and fatigue

	Stress – PSS-10 Score		Fatigue – Pichot Score	
	2-level	3-level	2-level	3-level
Intercept	16.7(0.23)	16.9 (0.38)	11.2 (0.25)	11.4 (0.5)
Random effects - $\sigma^2(\sigma)^*$				
Level 1 – Time	28.9 (5.3)	20.9 (4.6)	23.9 (4.8)	23.8 (4.9)
Level 2 – Healthcare worker	28.94 (5.3)	25.9 (5.1)	39.12 (6.3)	35.1 (6)
Level 3 – Ward		3.2 (1.8)		4.6 (2.1)
AIC ^a	15762.86	15737.24	15922.34	15896.45
ANOVA test p-value		< 2e-16		2.25 x 10e-07

^a AIC Akaike Information Criterion

Intra-class correlation coefficient (ICC) calculation

Using the 3-levels null models, ICC were computed, in order to quantify how much response variable variance is shared, or correlated, across different combinations of levels. ICC has been defined as “. . . an estimate of the expected (population) correlation between

two randomly chosen elements in the same group” [33]. As such, three different ICCs could be calculated to assess the influence of ward on change in level of stress and fatigue over time.

The level of stress and fatigue variance has been partitioned at all three levels. It can be easily shown that, the PSS-10 score and the Pichot score variations occurred due to temporal fluctuations (level 1, 42% for stress and 37.4% for fatigue), inter-individual heterogeneity (level 2, 52% for stress and 55.5% for fatigue) and ward-level specificities (level 3, 6% for stress and 7.2% for fatigue). In addition, these partitioned variances can be used to compute three different ICCs to assess the influence of ward on change in stress and fatigue level over time.

First, for the stress level, one level 3 ICC already estimated above (0.06), is interpreted as the expected correlation between two stress level drawn completely at random (from any time point), from two healthcare workers within the same service. Second, an alternative level 3 ICC estimate can be calculated as $(3.2 / (3.2+25.9) = 0.11\%)$ and interpreted as the expected correlation between the mean (i.e., averaged across all repeated measures) stress levels from two healthcare workers drawn completely at random from the same service. Finally, a level 2 ICC could also calculate by $(3.2 + 25.9 / (3.2+25.9+20.9) = 0.58)$ and is interpreted as the expected correlation between two repeated measurements sampled from the same healthcare workers. In the same way, 3 ICCs for fatigue are calculated and correspond to 0.07% for ICC level 3, 0.12 for the second ICC level 3 and 0.45 for the ICC level 2.

Supplementary table 5: Summary of missing values according to validated scales, by time of visits

	t0	t1	t2	t3	Total
PSS-10 score - Stress	7	3	5	7	22
Pichot score - Fatigue	1	0	2	0	3
Karasek score - Support from colleagues	6	4	1	1	12
Karasek score - Support from hierarchy	23	5	1	0	29
Siegrist score - Work overcommitment	1	1	0	1	3

In the whole sample (n =2472), high number of missing values were observed on support from the hierarchy and perceived stress, respectively 29 and 22 values. The lowest number of missing values were observed on fatigue and work overcommitment, all two 3 missing values. We count 12 missing values in the whole sample for the support from colleague’s variable. Before proceeding with the 3-levels analyses, missing values were imputed.